

TRK Fused Gene Rabbit Monoclonal Antibody

Catalog #: EAB22311

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IHC, IF/ICC	43	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:500-2000
IHC (Immunohistochemistry)	1:20-200
IF/ICC (Immunofluorescence/Immunocytochemistry)	1:20-200

Product Information

Conjugate	Unconjugate
Specificity	TRK Fused Gene Rabbit Monoclonal Antibody detects endogenous levels of TRK Fused Gene protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	Q92734
Entrez-Gene Id	10342

Product Description

Oncogenic rearrangements of the NTRK1 gene, which encodes the Trk A protein, are frequently detected in thyroid carcinomas. Such rearrangements fuse the NTRK1 tyrosine kinase domain to 5'-end sequences of different genes. TRK-T3 contains 1,412 nucleotides of NTRK1 preceded by 598 nucleotides belonging to TFG (TRK-fused gene), a ubiquitously expressed gene located on chromosome 3. The TRK-T3 protein within the TFG region contains a coiled-coil motif that gives the oncoprotein the capability to form complexes. The cytoplasmic TRK-T3 protein binds to and phosphorylates the Shc and SNT1/FRS2 adaptor proteins, both of which are involved in coupling the receptor tyrosine kinase to the mitogen-activated protein kinase pathway by recruiting Grb2/SOS. SHP-1 also interacts with and down-regulates TRK-T3.

For Reserch Use Only. Not For Use In Diagnostic Procedures