

# **Perilipin-2 Mouse Monoclonal Antibody**

# Catalog #: EAB22501

| Host/Isotype | Clonality  | Applications                 | MW (kDa) | Reactivity |
|--------------|------------|------------------------------|----------|------------|
| Mouse IgG1   | Monoclonal | WB, IHC-P, IF/ICC, FC, ELISA | 48       | Human      |

#### **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| WB(Western Blotting)                           | 1:500-2000   |
|--|--------------|
| IHC-P(Immunohistochemistry-Paraffin)           | 1:100-500    |
| IF/ICC(Immunofluorescence/Immunocytochemistry) | 1:100-500    |
| FC(Flow Cytometry)                             | 1:50-200     |
| ELISA(Enzyme-linked Immunosorbent Assay)       | 1:5000-20000 |

## **Product Information**

| Conjugate      | Unconjugate  |
|----------------|--|
| Specificity    | Perilipin-2 Mouse Monoclonal Antibody detects endogenous levels of Perilipin-2 protein .   |
| Purification   | Affinity purification  |
| Concentration  | 1mg/ml   |
| Format         | Liquid   |
| Formulation    | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.  |
| Shipping       | Gel Pack   |
| Storage        | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles.<br>Aliquots may be stored at +4°C for 1-2 weeks. |
| UniProt ID     | <u>Q99541</u>  |
| Entrez-Gene ID | <u>123</u>   |

## **Product Description**

Perilipin-2 also known as ADFP; ADRP belongs to the perilipin family, members of which coat intracellular lipid storage droplets. This protein is associated with the lipid globule surface membrane material, and maybe involved in development and maintenance of adipose tissue. However, it is not restricted to adipocytes as previously thought, but is found in a wide range of cultured cell lines, including fibroblasts, endothelial and epithelial cells, and tissues, such as lactating mammary gland, adrenal cortex, Sertoli and Leydig cells, and hepatocytes in alcoholic liver cirrhosis, suggesting that it may serve as a marker of lipid accumulation in diverse cell types and diseases. Alternatively spliced transcript variants have been found for this gene.

For Reserch Use Only. Not For Use In Diagnostic Procedures